

LIMITED prediction

These patients have a SAFE score less than 5 at 3 days post-stroke, and TMS cannot elicit MEPs in their paretic upper limb 3 – 7 days post-stroke. However, their total NIHSS score 3 days post-stroke was less than 7.

- This patient is most likely to have a limited upper limb outcome within the next 3 months.
- They can expect to regain some movement and possibly grasp function, though recovery of dextrous hand function is unlikely.
- Upper limb rehabilitation can focus on improving strength, active range of motion, and joint flexibility, and adapting daily activities to incorporate both upper limbs when necessary to achieve a task.
- This prediction is based on the patient’s current status, and is not a guarantee, as some people recover more or less than expected.

A version of this information has been given in written form and discussed with the patient (copy enclosed)

Signed:

Date:

LIMITED prediction

These patients have a SAFE score less than 5 at 3 days post-stroke, and TMS cannot elicit MEPs in their paretic upper limb 3 – 7 days post-stroke. However, their total NIHSS score 3 days post-stroke was less than 7.

- This patient is most likely to have a limited upper limb outcome within the next 3 months.
- They can expect to regain some movement and possibly grasp function, though recovery of dextrous hand function is unlikely.
- Upper limb rehabilitation can focus on improving strength, active range of motion, and joint flexibility, and adapting daily activities to incorporate both upper limbs when necessary to achieve a task.
- This prediction is based on the patient’s current status, and is not a guarantee, as some people recover more or less than expected.

A version of this information has been given in written form and discussed with the patient (copy enclosed)

Signed:

Date:

LIMITED prediction

These patients have a SAFE score less than 5 at 3 days post-stroke, and TMS cannot elicit MEPs in their paretic upper limb 3 – 7 days post-stroke. However, their total NIHSS score 3 days post-stroke was less than 7.

- This patient is most likely to have a limited upper limb outcome within the next 3 months.
- They can expect to regain some movement and possibly grasp function, though recovery of dextrous hand function is unlikely.
- Upper limb rehabilitation can focus on improving strength, active range of motion, and joint flexibility, and adapting daily activities to incorporate both upper limbs when necessary to achieve a task.
- This prediction is based on the patient’s current status, and is not a guarantee, as some people recover more or less than expected.

A version of this information has been given in written form and discussed with the patient (copy enclosed)

Signed:

Date:

LIMITED prediction

These patients have a SAFE score less than 5 at 3 days post-stroke, and TMS cannot elicit MEPs in their paretic upper limb 3 – 7 days post-stroke. However, their total NIHSS score 3 days post-stroke was less than 7.

- This patient is most likely to have a limited upper limb outcome within the next 3 months.
- They can expect to regain some movement and possibly grasp function, though recovery of dextrous hand function is unlikely.
- Upper limb rehabilitation can focus on improving strength, active range of motion, and joint flexibility, and adapting daily activities to incorporate both upper limbs when necessary to achieve a task.
- This prediction is based on the patient’s current status, and is not a guarantee, as some people recover more or less than expected.

A version of this information has been given in written form and discussed with the patient (copy enclosed)

Signed:

Date: