

TMS NOT APPROPRIATE

This patient's functional upper limb outcome cannot be predicted with SAFE scores alone.

Further investigation with Transcranial Magnetic Stimulation (TMS) and the use of NIHSS would be required to give an accurate prediction of this patient's upper limb recovery at 12 weeks, but this patient is not appropriate for TMS because:

TMS NOT APPROPRIATE

This patient's functional upper limb outcome cannot be predicted with SAFE scores alone.

Further investigation with Transcranial Magnetic Stimulation (TMS) and the use of NIHSS would be required to give an accurate prediction of this patient's upper limb recovery at 12 weeks, but this patient is not appropriate for TMS because:

TMS NOT APPROPRIATE

This patient's functional upper limb outcome cannot be predicted with SAFE scores alone.

Further investigation with Transcranial Magnetic Stimulation (TMS) and the use of NIHSS would be required to give an accurate prediction of this patient's upper limb recovery at 12 weeks, but this patient is not appropriate for TMS because:

TMS NOT APPROPRIATE

This patient's functional upper limb outcome cannot be predicted with SAFE scores alone.

Further investigation with Transcranial Magnetic Stimulation (TMS) and the use of NIHSS would be required to give an accurate prediction of this patient's upper limb recovery at 12 weeks, but this patient is not appropriate for TMS because:
